Acūtis

Instruction for STI specimen collection


Acutis Reveal™ STI

Collection for rectal swab specimens

Materials



Steps







Carefully insert swab into the rectum about 1-2 inches (3-5 cm) past the anal margin (the outside of the anus) and gently rotate swab clockwise for 5-10 seconds. Withdraw swab without touching skin.




4

While holding the swab in hand, unscrew the tube cap. Do not spill the tube contents.

Note: If the tube contents are spilled, discard and replace with a new Aptima Multitest Swab Specimen Collection Kit.



Immediately place the swab into the transport tube so black score line is at top of the tube.



Align the score line with the top edge of the tube and carefully break the shaft. Discard the top portion of the shaft.



Tightly screw cap onto the tube. The tube label provides a specimen source field for unique identification for specimen collection site location.



Sample type must be identified. Complete requisition using patient's name, date of birth, specimen collection date and time, and initials of the specimen collector for transport to laboratory.

Mailing instructions

- ightarrow Confirm patient name, DOB and collection date are legible on the vial
- ightarrow Place tube into the sealed compartment of the specimen bag
- ightarrow Be sure to include in the specimen bag the following items:
 - 1. Test requisition form with patient demographics
 - 2. Copy of front and back of insurance card
 - 3. Medical records indicating medical necessity
- ightarrow Place completed specimen bag(s) into the FedEx/UPS Lab Pak and seal the Lab Pak
- ightarrow Affix prepaid return label on the Lab Pak, and schedule a pickup
- → Send to Acutis Diagnostics: 400 Karin Lane, Hicksville, NY 11801 or 68 Culver Road, Suite 150B, Monmouth Junction, NJ 08852